K-State Wheat Demonstration Plots

Sherman County

Plot Location: 10 Mi North of Goodland

Plot Cooperator: F & J Farms

	Yield	Moisture	Test Weight	Protein
	(bu/A)	(%)	(%)	(%)
T-158	96.8	12.4	56.4	13.3
WB 4458	92.2	13.2	56.8	13.9
TAM 114	86.6	13.1	57.2	13.8
KanMark	85.0	11.7	57.0	13.6
Winterhawk	82.2	13.0	56.3	13.6
LCS Pistol	81.0	11.4	55.5	13.9
Denali/WB Grainfield/T	78.1	12.9	55.0	13.9
Oakley CL	72.1	12.0	54.0	15.2
Denali	68.6	12.1	54.2	13.5
Byrd/LCS Mint/Winterl	67.1	12.2	55.0	12.7
Byrd/LCS Mint/Wizard	65.6	12.2	54.8	13
TAM 112	65.2	12.8	56.0	13.8
WB Grainfield	59.1	12.8	52.4	14.6
Robidoux	57.7	12.3	54.4	13.5
LCS Mint	50.1	12.0	53.4	13.5
LCS Wizard	44.2	11.8	54.1	13.6
Judee	41.9	11.1	52.0	15.7
Quake	35.0	10.5	51.3	15.6
TAM 113	29.7	11.7	54.0	14.6
Cowboy	24.0	11.0	53.0	13.6
Hatcher	4.8	11.0	54.4	13.8
Average	61.3	12.1	54.6	13.9

^{*}Aborted: The 3 varieties on the north side of the plots had no data due to winter injury because of open north exposure. Alphabetical: Antero, Brawl CL, Byrd

October 6, 2015

Drilled: at 71 lbs per acre

with plot size 750' x 30 ft = 0.52 ac/plot with starter fertilizer: 14 N and 36 P

Sprayed for stripe rust early in infection process.

Harvested July 20, 2015

These results are from demonstration plots and should be used with replicated performance test data. AgriPro results are not included in the plot results, per AgriPro's wishes.

If you have any questions, please contact K-State Agronomist, Jeanne Falk Jones at (785) 443-3403.